

Weitere Bestandteile

	Allurarot	Tartrazin	Gelborange S	Titandioxid	Indigocarmin	Eisen(II,III)-oxidhydrat	Eisen(III)-oxidhydrat	Eisen(III)-oxidhydrat,schwarz	Drucktinte, propylenglycolhaltig	Drucktinte	Gelatine
Bristol Meyers Squibb				x	x			x	x		
123-Acurae Pharma *1				x	x					x	
ABZ-Pharma *2				x	x		x		x		x
Accord											
Aluid			x	x		x	x	x			x
Aristo	x	x	x	x					x		x
Basics				x	x			x		x	x
beta				x	x			x	x		
Cipla Europe NV				x	x			x	x		
Devatis				x	x			x	x		
Glenmark				x	x			x	x		
Heumann	x	x	x	x					x		x
Hexal				x	x			x	x		
Mylan/Viatrix				x					x		x
Ohre Pharma/Stragen				x	x			x	x		x
Piramal	x	x	x	x					x		x
Puren					x		x	x			
Ratiopharm *2				x	x		x		x		x
Stada	x	x						x			
TAD *1				x	x			x	x		
Zentiva				x	x			x	x		

*1 Individuelle sonstige Bestandteile: Mannitol, Cellulose, Maisstärke, Weinsäure, Glycerol, Hypromellose, Carrageen

*2 Individuelle sonstige Bestandteile: Siliciumdioxid, Cellulose, Croscarmellose, Talkum

Verfasser: Mathias Nietzke, Apotheker Leiter Zytostatika-Abteilung des St.-Johannes Hospitals Dortmund Mathias.nietzke@joho-dortmund.de

Quellenangabe: <https://www.myelom.org/grundlagen/wechselwirkungen-von-arzneimitteln.html>